AMENDMENTS TO THE CLAIMS: This listing of claims replaces all prior versions and listings of claims in the instant patent application.

Listing of claims:

- 1. (currently amended) A method of detecting Graves' disease or rheumatoid arthritis in a patient comprising
- (a) obtaining a biological sample comprising fibroblasts from the patient, and
- (b) <u>detecting in said biological sample</u> <u>measuring</u> the <u>activation of fibroblasts</u> by binding of disease specific IgG <u>with to the IGF-1 receptor</u> (IGF-1R) relative to a control wherein <u>presence of IgG-activated fibroblasts compared</u> <u>an elevated level of IgG IGF-1 biding relative</u> to the control indicates Graves' disease or <u>rheumatoid arthritis</u>.
- 2. (cancelled)
- 3. (currently amended) The method of claim 1 wherein the determination detecting is accomplished by measures measuring the level of a chemical marker expressed by said IgG-activated T cells fibroblasts in said biological sample, wherein an elevated level of the marker compared to the control indicates presence of said IgG-activated fibroblasts.
- 4. (original) The method of claim 3 wherein the marker is RANTES.
- 5. (currently amended) The method of claim 3 wherein the marker is IC-16 IL-16.
- 6. (currently amended) The method of claim 2 wherein the determination is detecting is accomplished by exposing T-cells to said biological sample comprising fibroblasts and measuring T-cell migration toward said fibroblasts, wherein an increase in the migration of said fibroblasts relative to the control indicates presence of said IgG-activated fibroblasts.
- 7. (original) The method of claim 1 wherein the patient is human.
- 8. (currently amended) The method of claim 1 wherein the biological sample is selected from a group consisting of: blood, urine, synovial fluid, ascites, <u>and</u> tissues.
- 9. (new) A method of detecting the presence of antibody-activated fibroblasts, said method comprising
- (a) obtaining a biological sample comprising fibroblasts from the patient;

10/038,509

- (b) contacting said sample with an antibody specific for IL-16
- (c) detecting the level of IL-16 released by said fibroblasts relative to a control, wherein an elevated level of IL-16 detects the presence of antibody-activated fibroblasts.
- 10. (new) A method of detecting the presence of antibody-activated fibroblasts, said method comprising
- (a) obtaining a biological sample comprising fibroblasts from the patient;
- (b) contacting said sample with an antibody specific for RANTES;
- (c) detecting the level of RANTES released by said fibroblasts relative to a control, wherein an elevated level of RANES detects the presence of antibody-activated fibroblasts.
- 11. (new) A method of detecting the presence of antibody-activated fibroblasts, said method comprising
- (a) obtaining a biological sample comprising fibroblasts from the patient;
- (b) contacting said sample with antibodies specific for IL-16 and RANTES;
- (c) detecting the levels of IL-16 and RANTES released by said fibroblasts relative to a control, wherein an elevated level of both IL-16 and RANTES detects the presence of antibody-activated fibroblasts.